



Introduction to telescopes  
and types of instruments III

*the Very Large Telescope Interferometer*  
*-VLTI-*

---

*Claudia Paladini*


ESO Operations Staff Astronomer

# Why Interferometry?

Objects	Wavelength	Diameter	Telescope diameter
Circumstellar envelope around evolved star	11 $\mu\text{m}$	50 mas	45 m (more than the ELT!)
Vulcan on a Jupiter satellite	5 $\mu\text{m}$	10 mas	100 m
Nucleus of AGN	2.2 $\mu\text{m}$	< 1 mas	> 400 m
Spot on the photosphere of a solar-type star	0.5 $\mu\text{m}$	0.07 mas	1500 m

# Principles of Interferometry

- Not a single dish, but light combined from 2,3,4... telescopes
- **Gain:** angular resolution
- **Cost:** sensitivity





*«The complex visibility is the Fourier transform of the source intensity distribution on the sky at the spatial frequencies corresponding to the projected baseline on the sky per observing wavelength.»*


---

**Van-Cittert-Zernicke theorem**


**Linking the complex visibility**

**to the intensity distribution of the object in the sky**


# Van-Cittert-Zernicke theorem



Choose your preferred object



Project the baseline on your object and collapse the intensity in that direction




Take 1-D Fourier Transform

# What is the observable?

We observe **FRINGES** and we measure a complex quantity called **VISIBILITY**

- fringe visibility is the contrast between fringes => **Angular dimension of the object.**
- fringe phase related to the location of fringes. => **Symmetry of the object**



Interferometric Fringes from Star with Different Angular Diameters  
(Simulation)

ESO PR Photo 10d/01 (18 March 2001)

© European Southern Observatory

*left: “real star”*

*center: star observed by single dish*


*right: star observed by interferometer*

# The u-v plane

The **complex visibility** is a function of the **spatial frequencies or u-v coordinates**


$$u = \frac{B_x}{\lambda} \quad \frac{B_p}{\lambda} = \sqrt{u^2 + v^2}$$

$$v = \frac{B_y}{\lambda}$$



- Projected baseline (baseline as seen from the star) is what matters
- Sidereal motion changes this projection
- Changing the wavelength also changes the spatial frequency

# Baseline effect



Interferometric Fringes at Different Telescope Baselines  
(Simulation)

ESO PR Photo 10e/01 (18 March 2001)

© European Southern Observatory



Short baseline  
Long baseline



higher contrast  
smaller contrast ( $V \sim 0$ )

# About the phase: closure phase

- In visible & infrared we cannot measure the “pure” phase because of the atmosphere variation


$$\Phi_{1,2} = \phi_{1,2}^{\text{obj}} + \phi_2^{\text{atm}} - \phi_1^{\text{atm}}$$

$$\Phi_{2,3} = \phi_{2,3}^{\text{obj}} + \phi_3^{\text{atm}} - \phi_2^{\text{atm}}$$

$$\Phi_{3,1} = \phi_{3,1}^{\text{obj}} + \phi_3^{\text{atm}} - \phi_1^{\text{atm}}$$

- In optical near-infrared interferometry we use the “**closure phase**”

Credit: Monnier+2006



$$\Psi_{1,2,3} = \phi_{1,2}^{\text{obj}} + \phi_{2,3}^{\text{obj}} + \phi_{3,1}^{\text{obj}}$$

# About the Phase: differential phase

Definition:

*Choose a reference spectral channel.*

*The differential phase is the phase difference between different spectral channels.*




If you have only 2 telescopes and few wavelength channels:  
**differential phase.**




The differential phase is sensitive to the astrometric position


# Modeling: Uniform disc



Software: ASPRO




U1-U2-U3  
 Wavelength 2,000  $\mu$   
 Declination  $-14.8^\circ$   
 Model: C DISK





# Modeling: binary


Binary stars:  
 diameter of the primary resolved  
 star 10 mas  
 Flux ratio = 0.8  
 $\theta = 45$  degree





Software: ASPRO




# Imaging



Credit: Paladini++2018




# Imaging



Credit: Paladini++2018


# Some applications



# Observations in practice

Between us and the star there is atmosphere and the instrument

We measure visibility as well as **instrumental effect & atmosphere (transfer function TF)**



# Calibration of the visibility

To correct for these effects, we observe a source with a known diameter  $\theta$ , the calibrator

$$(1) \quad \mu_{obs}^2 = \frac{V_{intrinsic}^2}{T^2}$$

For the calibrator, assuming a Uniform disk distribution:

$$(2) \quad V_{intrinsic}(u, v) = 2 \frac{J_1(\pi\theta r)}{\pi\theta r}$$


Combining (1) and (2) one can derive the transfer function T and calibrate the intrinsic visibility of the  
SCIENCE

# Example of calibration at VLTI

*PIONIER calibration sequences are CAL-SCI-CAL or CAL-SCI-CAL-SCI-CAL*

Seeing: ASM  
vs. constraints

Closure phases



Visibilities

Science target:  
orange star

Calibrator:  
gray dot &  
blue cross







## VLTI acts like a virtual 200 m telescope

- Combining 4UTs (8m telescope), maximum baseline 130 m, resolution  $\sim 2$  mas at 2 micron
- Combining 4ATs (1.8m telescope), maximum baseline 200 m (*from October 2023*), resolution  $\sim 0.8$  mas at 1.5 micron



La Silla Observing School, Chile, February 2025





# VLT today

## Auxiliary Telescopes equipped with NAOMI Adaptive Optics

- R ~ 15 mag in excellent conditions (visitor mode)
- R  $\leq$  12.5 in standard conditions (service mode)

## Unit Telescopes equipped with

- GPAO + Natural guide star Adaptive Optics V = 12.5
- GPAO + Laser Guide star from 2026 will improve limits by 5 mag
- CIAO Adaptive Optics GRAVITY only, K  $\leq$  10

# VLTI today

## PIONIER

- H band ( $\lambda \sim 1.6\mu\text{m}$ )
- R~50
- ATs limit H ~ 9 mag
- Not used on the UTs (injection)

## GRAVITY

- K band ( $\lambda \sim 2.2\mu\text{m}$ ),
- R~20, 500 and 4000
- Fringe tracker (up to 2" off-axis)

## MATISSE

- L,M,N bands ( $\lambda \sim 3$  to  $12\mu\text{m}$ ),
- R~30, 500, 1000 and 3500
- GRAVITY as a fringe tracker



**KEEP  
CALM  
AND  
OBSERVE  
WITH  
PIONIER**













Image credit: GRAVITY consortium

# Observing the Universe in Motion: 5 years of GRAVITY





19+ mag limiting magnitude & polarimetry




2 x 4 milli-arcsec resolution imaging


Micro-arcsec spectral differential astrometry




High resolution spectroscopy

<50 μas imaging astrometry





Betelgeuse's Great Dimming Event in high resolution  
Drevon et al. 2023




A close-up view of Messier  
77's active galactic nucleus  
Gómez-Rosas et al. 2022



# MATISSE

# VLTI tomorrow: going faint with GRAVITY+




+5mag and better sky coverage:

- off-axis fringe tracking (2" → 30")
- Laser Guide Star on every UT
- Higher performance Adaptive Optics
- Better vibration control

Science Cases (Kmag~22):

- The Galactic Centre
- Galaxy AGN coevolution and the masses of supermassive Black Holes (up to  $z \sim 2.5$ )
- Characterization of exoplanets
- Young suns and their planet-forming disks



# Thank you!

---

**Claudia Paladini**  
**cpaladin@eso.org**

-  @ESO Astronomy
-  @eso astronomy
-  @ESO
-  european-southern-observatory
-  @ESOobservatory

