

Problem 3: One-dim. UIR of the Braid group B_n

Generators obey:

$$\varepsilon_i \varepsilon_j = \varepsilon_j \varepsilon_i \quad \text{for } |i - j| > 1$$

$$\varepsilon_i \varepsilon_{i+1} \varepsilon_i = \varepsilon_{i+1} \varepsilon_i \varepsilon_{i+1}$$

a) Make ansatz for 1-D UIR for generators: $D^\alpha(\varepsilon_j) = e^{i\alpha_j}$

- Unitarity: $\alpha_j \in [0, 2\pi[$
- First property: $D_i^\alpha(\varepsilon_i \varepsilon_j) = e^{i\alpha_i + i\alpha_j} = D^\alpha(\varepsilon_i) D^\alpha(\varepsilon_j)$
- Second property: $D^\alpha(\varepsilon_i \varepsilon_{i+1} \varepsilon_i) = e^{i(\alpha_i + \alpha_{i+1} + \alpha_i)} \stackrel{!}{=} D^\alpha(\varepsilon_{i+1} \varepsilon_i \varepsilon_{i+1}) = e^{i(\alpha_{i+1} + \alpha_i + \alpha_{i+1})}$
 $\implies \alpha_{i+1} = \alpha_i \bmod 2\pi \implies \alpha_{i+1} = \alpha_i \text{ for all } i.$

Hence $\alpha \in [0, 2\pi[$ characterises the full set of all 1-dim. UIR of B_n

b) Restriction to S_n implies a third property $\varepsilon_i^2 = e$

$$\implies D^\alpha(\varepsilon_i^2) = e^{2i\alpha} \stackrel{!}{=} 1 \implies \alpha = \begin{cases} 0 & \text{trivial reps} \\ \pi & \text{anti-sym. reps} \end{cases}$$

Comments:

Consider a quantum system of n identical particles characterised by the n -particle wave function $\Psi(x_1, \dots, x_n)$. The permutation of these particles may be represented by a permutation of the positions x_i : $P = \begin{pmatrix} 1 & 2 & \cdots & n \\ \pi_1 & \pi_2 & \cdots & \pi_n \end{pmatrix}$. The representation of such a permutation in the n -particle $L^2(\mathbb{R}^{3n})$ is given by

$$D^\alpha(P)\Psi(x_1, \dots, x_n) := \Psi(x_{\pi_1}, \dots, x_{\pi_n})$$

For identical particle physics should not change, that is, $|\Psi(x_1, \dots, x_n)|^2 = |\Psi(x_{\pi_1}, \dots, x_{\pi_n})|^2$.

Hence, D^α must be a 1-dim. UIR of S_n :

- Trivial reps.: $D^0(P)\Psi(x_1, \dots, x_n) := \Psi(x_1, \dots, x_n) = \Psi(x_{\pi_1}, \dots, x_{\pi_n})$. **Bosons**
- Anti-sym. reps.: $D^\pi(P)\Psi(x_1, \dots, x_n) := (-1)^k \Psi(x_1, \dots, x_n)$. **Fermions**
 $k = \text{number of transposition in } P$

For elementary particles these are the only physical 1D UIR of B_n . However, for "quasi particles" (collective excitation in a solid) any α may be realized. These are called **Anyons**. See, for example,

J. Jacak, R. Gonczarek, L. Jacak and I. Józwiak, *Application of Braid Groups in 2D Hall System Physics* (World Scientific, Singapore, 2012) <https://doi.org/10.1142/8512>

Problem 4: One-dim. UIR of $SO(2) \simeq U(1)$

Defining Reps of $SO(2)$ in \mathbb{R}^2

$$g(\varphi) = \begin{pmatrix} \cos \varphi & -\sin \varphi \\ \sin \varphi & \cos \varphi \end{pmatrix}, \quad \varphi \in [0, 2\pi[.$$

a) Consider the homomorphism

$$H : \begin{array}{ccc} SO(2) & \rightarrow & U(1) \\ g(\varphi) & \mapsto & e^{i\varphi} \end{array}$$

Note that using the trigonometric addition theorems we have

$$g(\varphi_1)g(\varphi_2) = \begin{pmatrix} \cos \varphi_1 & -\sin \varphi_1 \\ \sin \varphi_1 & \cos \varphi_1 \end{pmatrix} \begin{pmatrix} \cos \varphi_2 & -\sin \varphi_2 \\ \sin \varphi_2 & \cos \varphi_2 \end{pmatrix} = \begin{pmatrix} \cos(\varphi_1 + \varphi_2) & -\sin(\varphi_1 + \varphi_2) \\ \sin(\varphi_1 + \varphi_2) & \cos(\varphi_1 + \varphi_2) \end{pmatrix}$$

Hence, $g(\varphi_1)g(\varphi_2) \mapsto e^{i(\varphi_1 + \varphi_2)}$.

Obviously this is invertable and H is actually an isomorphism.

Alternatively, let $\vec{x} = (x_1, x_2)^T \in \mathbb{R}^2$, then we can map it onto $z := x_1 + ix_2 \in \mathbb{C}$. Hence the rotation $g(\varphi)$ in \mathbb{R}^2 is replaced by the multiplication with phase $e^{i\varphi}$ in \mathbb{C} .

Consider unitary transformation

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}, \quad U^\dagger = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -i \\ -i & 1 \end{pmatrix} = U^{-1},$$

then the 2-dim. matrix reps. is reduced to

$$U^\dagger g(\varphi) U = \begin{pmatrix} e^{i\varphi} & 0 \\ 0 & e^{-i\varphi} \end{pmatrix}.$$

b) Ansatz for 1-dim. UIR: $D^\alpha(g) = e^{i\alpha(\varphi)}$ with $\alpha(\varphi) \in [0, 2\pi]$.

$$\bullet \quad g(\varphi_1)g(\varphi_2) = g(\varphi_1 + \varphi_2) \implies \alpha(\varphi_1) + \alpha(\varphi_2) = \alpha(\varphi_1 + \varphi_2) \bmod 2\pi.$$

Hence α is linear in φ .

$$\bullet \quad g(0) = \mathbf{1} = g(2\pi) \implies \alpha(0) = 2\pi m \text{ and } \alpha(2\pi) = 2\pi n \text{ with } m, n \in \mathbb{Z}$$

Conclusion: $\alpha(\varphi) = m \cdot \varphi$ and

$$\boxed{D^m(g) = e^{im\varphi} \quad \text{with} \quad m \in \mathbb{Z}}$$

is 1-dim. UIR of $SO(2) \simeq U(1)$.

Problem 5: Translations in \mathbb{R}^3 and its 1D UIR

$$T^3 : \begin{cases} \mathbb{R}^3 \rightarrow \mathbb{R}^3 \\ \vec{a} \mapsto \vec{a} + \vec{x} \end{cases} \quad \text{with group element} \quad g(\vec{x}) = \begin{pmatrix} \mathbf{1}_3 & \vec{x} \\ \vec{0}^T & 1 \end{pmatrix} \quad \text{and} \quad \vec{x} \in \mathbb{R}^3$$

a) Neutral element $g(\vec{0}) = \mathbf{1}$ obvious.

Hence we only need to verify the group law of translations, which implies $g(\vec{x})g(\vec{y}) = g(\vec{x} + \vec{y})$:

$$g(\vec{x})g(\vec{y}) = \begin{pmatrix} \mathbf{1}_3 & \vec{x} \\ \vec{0}^T & 1 \end{pmatrix} \begin{pmatrix} \mathbf{1}_3 & \vec{y} \\ \vec{0}^T & 1 \end{pmatrix} = \begin{pmatrix} \mathbf{1}_3 & \vec{x} + \vec{y} \\ \vec{0}^T & 1 \end{pmatrix} = g(\vec{x} + \vec{y})$$

Obviously the inverse of $g(\vec{x})$ is given by $g(-\vec{x})$ and T^3 is abelian.

However,

$$g^{-1}(\vec{x}) = g(-\vec{x}) = \begin{pmatrix} \mathbf{1}_3 & -\vec{x} \\ \vec{0}^T & 1 \end{pmatrix} \neq g^\dagger(\vec{x}) = \begin{pmatrix} \mathbf{1}_3 & \vec{0} \\ \vec{x}^T & 1 \end{pmatrix} \notin T^3$$

The above matrix representation of T^3 is neither unitary nor irreducible. It acts on \mathbb{R}^4 with $(0, 0, 0, 1)^T$ spanning an invariant subspace.

b) Ansatz $D_{\vec{k}}(g(\vec{x})) = e^{i\vec{k} \cdot \vec{x}}$ is representation as

$$D_{\vec{k}}(g(\vec{0})) = 1 \quad \text{and} \quad D_{\vec{k}}(g(\vec{x} + \vec{y})) = D_{\vec{k}}(g(\vec{x}))D_{\vec{k}}(g(\vec{y}))$$

Unitarity follows from $D_{\vec{k}}(g(-\vec{x})) = D_{\vec{k}}^*(g(\vec{x}))$